



Influence of cerium precursors on the structure and reducibility of mesoporous CuO-CeO₂ catalysts for CO oxidation

Lei Qi^a, Qiang Yu^a, Yue Dai^a, Changjin Tang^a, Lianjun Liu^a, Hongliang Zhang^a, Fei Gao^{b,**}, Lin Dong^{a,b,*}, Yi Chen^a

^a Key Laboratory of Mesoscopic Chemistry of MOE, School of Chemistry and Chemical Engineering, Nanjing University, Nanjing 210093, PR China

^b Jiangsu Key Laboratory of Vehicle Emissions Control, Center of Modern Analysis, Nanjing University, Nanjing 210093, PR China

ARTICLE INFO

Article history:

Received 30 November 2011

Received in revised form 22 February 2012

Accepted 25 February 2012

Available online 5 March 2012

Keywords:

Cerium precursor

Synergistic effect

Reducibility

CuO-CeO₂

CO oxidation

ABSTRACT

This work investigated the effects of cerium precursors [Ce(NO₃)₃ and (NH₄)₂Ce(NO₃)₆] on the structure, surface state, reducibility and CO oxidation activity of mesoporous CuO-CeO₂ catalysts. The catalysts were characterized by TG-DTA, XRD, LRS, N₂ adsorption-desorption, HRTEM, XPS, H₂-TPR and in situ FT-IR. The obtained results suggested that the precursors exerted a great influence on the properties of CuO-CeO₂ catalysts: (1) compared with the catalysts from Ce(III) precursor, the derived Ce(IV) precursor catalysts showed smaller grain size, higher BET surface area, narrower pore size distribution, whereas their reducibility and activities were not enhanced. (2) In contrast, the catalysts from Ce(III) precursor without excellent texture displayed high reducibility and activities for CO oxidation due to the high content of Ce³⁺, following the redox equilibrium of Cu²⁺ + Ce³⁺ ⇌ Cu⁺ + Ce⁴⁺ shifting to right to form more stable Cu⁺ species, which was the origin of synergistic effect. The synergistic effect between copper and cerium was the predominant contributor to the improved catalytic activities of CuO-CeO₂ catalysts, instead of structural properties.

© 2012 Elsevier B.V. All rights reserved.

1. Introduction

In recent years, ceria (CeO₂) has been received increasing attentions and found tremendous applications in environmental catalysis due to its high oxygen storage-release capacity associated with Ce⁴⁺/Ce³⁺ redox cycle [1,2]. It was well documented that the catalytic activities of ceria-based catalysts were greatly promoted not only by precious metals, but also by base metals, particularly copper species [3–6]. For instance, CuO-CeO₂ catalysts had been shown to be very active for CO oxidation, exhibiting specific activities comparable with or even superior to precious metal catalysts [7,8].

In principle, the high activity of CuO-CeO₂ is correlated with the synergism of copper-ceria interactions, which facilitates the electron exchanges between Cu²⁺/Cu⁺ and Ce³⁺/Ce⁴⁺, with both components being significantly more readily reduced or oxidized than the corresponding independent ones [9–11]. The redox cycles between Cu²⁺ and Cu⁺ (and concomitantly Ce⁴⁺ and Ce³⁺) are often regarded as the key steps of the reaction mechanism of CO oxidation

and consequently, as well as in determining the catalytic behavior [12]. Thus, the CuO-CeO₂ catalysts with facile electron exchanges can easily form Cu⁺ species for CO adsorption, which is an important factor to promote activity of CO oxidation [13]. Another important factor is the presence of the oxygen vacancy provided by CeO₂, which can in turn generate active oxygen for the reaction [14]. Other than these, unique texture of ceria-based catalysts is also advantageous for their catalytic performances. For example, mesoporous CuO-CeO₂ catalysts had been reported to display superior activity toward CO oxidation in comparison with usually used CuO/CeO₂ catalysts [15–17]. Consequently, controllable synthesis of mesoporous CuO-CeO₂ catalysts, which possess facile redox process and a high concentration of oxygen vacancies, will be beneficial to the enhancement of activity toward CO oxidation.

Until now, Ce(NO₃)₃ and (NH₄)₂Ce(NO₃)₆ are two of the most widely used precursors to synthesize CuO-CeO₂ catalysts. Tang et al. [18] prepared CuO-CeO₂ catalysts by precipitation method and used Ce(NO₃)₃ as cerium precursor. They pointed out that the discrepancies in the dispersion of copper species and the degree of interaction between copper species and ceria determined the reversible redox properties, and consequently the catalytic performance of CO oxidation. Li et al. [19] obtained CuO-CeO₂ catalysts from (NH₄)₂Ce(NO₃)₆ by urea co-precipitation-gelation method. They found that the reducibility of ceria was promoted by the addition of copper, leading to the significantly enhanced activity in the water-gas shift reaction (WGS). However, to our knowledge,

* Corresponding author at: School of Chemistry and Chemical Engineering, Nanjing University, Hankou Road 22#, Nanjing, Jiangsu 210093, China.

Tel.: +86 25 83592290; fax: +86 25 83317761.

** Corresponding author. Tel.: +86 25 83592290; fax: +86 25 83317761.

E-mail addresses: gaofei@nju.edu.cn (F. Gao), donglin@nju.edu.cn (L. Dong).




Fig. 1. TG-DTA curves of the uncalcined samples: (a) Ce(III)Cu_x; (b) Ce(IV)Cu_x.

little attention has been paid to comparing the effect of cerium precursors on the structure, surface state, redox property and CO oxidation activity of mesoporous CuO-CeO₂ catalysts. On the other hand, previous researches proved that many factors could influence the activity of CuO-CeO₂ catalysts, such as surface area, porous structure [15], synergistic effect between copper and cerium [12], oxygen vacancy [14], etc. Nevertheless, the contributions of these factors to the catalytic activity and the predominant contributor to the enhanced activity are still not clear.

In this work, a series of mesoporous CuO-CeO₂ catalysts were prepared by a surfactant-assisted co-precipitation method and characterized by means of TG-DTA, XRD, LRS, N₂ adsorption-desorption, HRTEM, H₂-TPR, in situ FT-IR and CO oxidation model reaction. The studies have mainly focused on (1) exploring the influence of cerium precursors (i.e. Ce(NO₃)₃ or (NH₄)₂Ce(NO₃)₆) on the structure, surface state, reduction property and CO oxidation activity of CuO-CeO₂ catalysts; (2) investigating the contribution of structural property and synergistic effect to the reducibility and catalytic activity of CO oxidation, and deciding the major contributor to improve the activity.

2. Experimental

2.1. Catalyst preparation

All chemicals were of analytical grade and used as received without further purification. CuO-CeO₂ catalysts with different Cu contents were synthesized by a surfactant-assisted co-precipitation method. In a typical synthesis procedure, 6 mmol cetyltrimethyl ammonium bromide (CTAB) was dissolved in 200 mL distilled water at room temperature, followed by the addition of 10 mmol Ce(NO₃)₃·6H₂O or (NH₄)₂Ce(NO₃)₆ and a calculated amount of Cu(NO₃)₂·3H₂O. After that, an aqueous solution of NaOH (0.2 mol L⁻¹) was added dropwise to the pre-mixed solution under vigorous stirring until pH 9. The obtained suspension was aged at 90 °C for 3 h. The precipitate was filtered, washed and dried at 110 °C for 12 h. Subsequently, the resultant powders were calcined at 400 °C for 5 h in air. The catalysts were denoted as Ce(III)Cu_x or Ce(IV)Cu_x, where x stands for the percentage of Cu/(Cu + Ce) molar ratio. The nominal contents of Cu were 0, 5, 15, 30 and 50 mol%, respectively.

Table 1

Chemical composition, grain size, lattice parameter, full width at half maximum (FWHM) of F_{2g} vibration band, the area ratio of Raman bands at 590 and 460 cm^{-1} , surface area, pore size, pore volume and H_2 consumption of CuO-CeO_2 catalysts.

Catalysts	Cu (mol%) ^a	Cu/Ce ^b	Cu/Ce ^c	D (nm) ^d	a (nm) ^e	FWHM (cm^{-1}) ^f	A_{590}/A_{460} ^g	S_{BET} ($\text{m}^2 \text{g}^{-1}$) ^h	r (nm) ⁱ	V ($\text{cm}^3 \text{g}^{-1}$) ^j	Theoretical H_2 consumption ($\mu\text{mol g}^{-1}$)	Actual H_2 consumption ($\mu\text{mol g}^{-1}$)
Ce(III)Cu0	0	0	0	9.5	0.5427	33	0.05	143	8.37	0.30	–	–
Ce(III)Cu5	5.51	0.06	0.20	8.1	0.5423	35	0.22	130	7.20	0.23	327.6	789.3
Ce(III)Cu15	12.99	0.15	0.33	7.8	0.5418	50	0.23	149	8.59	0.32	835.4	1397.0
Ce(III)Cu30	28.10	0.39	0.64	6.6	0.5414	47	0.15	137	11.88	0.41	1874.9	2219.6
Ce(III)Cu50	51.63	1.07	0.95	5.2	0.5424	41	0.16	123	9.73	0.30	4140.4	4152.1
Ce(IV)Cu0	0	0	0	5.0	0.5431	32	0.02	177	4.10	0.18	–	–
Ce(IV)Cu5	5.68	0.06	0.15	4.2	0.5428	37	0.04	218	3.72	0.20	337.7	1000.3
Ce(IV)Cu15	14.31	0.17	0.34	4.1	0.5430	52	0.14	214	3.73	0.20	892.1	1407.7
Ce(IV)Cu30	31.58	0.46	0.62	3.9	0.5392	47	0.10	179	3.89	0.17	2200.0	2383.0
Ce(IV)Cu50	52.08	1.09	0.94	3.7	0.5423	41	0.11	166	5.80	0.24	4170.6	4188.0

^a Actual Cu content (mol%) determined by XRF.

^b Actual atomic ratio (Cu/Ce) determined by XRF.

^c Surface atomic ratio (Cu/Ce) determined by XPS.

^d Grain size of CeO_2 determined from the XRD diffraction peak by Scherrer equation.

^e Lattice parameter calculated from characteristic XRD peaks of CeO_2 by Bragg's law.

^f FWHM calculated from the characteristic peak of CeO_2 (F_{2g}) in Raman spectra.

^g The area ratio of Raman bands at 590 and 460 cm^{-1} .

^h Surface area determined from BET equation.

ⁱ Average pore radius determined from BJH method.

^j Pore volume.

2.2. Catalyst characterization

Thermogravimetry and differential thermal analysis (TG–DTA) of the catalysts was carried out on a Netzsch thermoanalyzer STA 449C with a heating rate of $10^\circ\text{C min}^{-1}$ in a flowing air.

The actual Cu contents in these CuO-CeO_2 catalysts were analyzed by X-ray fluorescence spectrometer (ARL-9800).

X-ray diffraction (XRD) patterns were recorded on a Philips X'pert Pro diffractometer using Ni-filtered $\text{Cu K}\alpha$ radiation (0.15418 nm). The X-ray tube was operated at 40 kV and 40 mA.

Laser Raman spectroscopy (LRS) analysis was performed on a Jobin-Yvon (France-Japan) T64000 type Laser Raman spectroscopy using Ar^+ laser beam. The Raman spectra were recorded with an excitation wavelength of 514 nm and the laser power of 300 mW.

The surface areas of the catalysts were determined via nitrogen adsorption at -196°C with the Brunauer–Emmet–Teller (BET) method using a Micrometrics ASAP-2020 adsorption apparatus. The pore size distribution was calculated from the desorption branch of the N_2 isotherm by using the Barrett–Joyner–Halenda (BJH) algorithm. Prior to each measurement, the sample was pretreated at 300°C for 4 h under vacuum.

High resolution transmission electron microscope (HRTEM) investigations were carried out using a JEM-2100 instrument at an acceleration voltage of 200 kV.

X-ray photoelectron spectroscopy (XPS) analysis was performed on a PHI5000 VersaProbe high performance electron spectrometer, using monochromatic $\text{Al K}\alpha$ radiation (1486.6 eV) operating at an accelerating voltage of 15 kV. All binding energies (BE) were referenced to the adventitious C1s at 284.6 eV. This reference gave BE values with an accuracy at ± 0.1 eV.

H_2 temperature-programmed reduction (H_2 -TPR) experiments were carried out in a quartz U-tube reactor connected to a thermal conduction detector with 7.3% H_2/Ar as reductant. 50 mg sample was used for each measurement. Before reduction, the sample was pretreated in a N_2 stream at 200°C for 1 h and then cooled to room temperature. After that, H_2/Ar mixture was switched on and the sample was heated with a heating rate of $10^\circ\text{C min}^{-1}$. The overlapped peaks were deconvoluted fitting with Gaussian–Lorentzian curves using the attached tool of origin PFM (Peak Fitting Module).

In situ Fourier transform infrared spectroscopy (FT-IR) was collected from 1000 to 4000 cm^{-1} with a resolution 4 cm^{-1} on a Nicolet

5700 FT-IR instrument (Thermo Electron Corporation, USA), using 64 scans per spectrum. The CuO-CeO_2 catalyst (about 10 mg) was pressed into a thin self-supporting wafer and mounted inside a high temperature cell (HTC-3, Harrick Scientific Corporation, USA). The wafer was pretreated for 1 h at 200°C in the flowing N_2 atmosphere. After cooled to room temperature, the sample was conducted to a controlled stream of 2% CO/N_2 or a mixture of 2% CO/N_2 and pure O_2 for 30 min. In situ CO adsorption and CO and O_2 co-adsorption FT-IR spectra were recorded at various target temperatures with subtraction of the corresponding background reference.

2.3. Catalytic activity measurements

The CO oxidation activities of the catalysts were measured in a flow micro-reactor with a gas composition of 1.6 vol% CO, 20.8 vol% O_2 and 77.6 vol% N_2 at a space velocity of $30,000 \text{ mL g}^{-1} \text{ h}^{-1}$, and 50 mg catalyst was used for each measurement. The catalyst was pretreated in a N_2 stream at 200°C for 1 h and then cooled to room temperature, after that, the mixed gases were switched on. Two columns and thermal conductivity detector (TCD) were used for the purpose of analyzing the production, column A with $13\times$ molecular sieve for separating O_2 , N_2 and CO, and column B, packed with Porapak Q for separating CO_2 .

3. Results and discussion

3.1. TG–DTA results

As shown in the TG–DTA curves (Fig. 1), the processes of weight loss are similar and can be divided into two steps for all samples. The initial weight loss below 180°C (denoted as step I) can be attributed to the desorption of water and condensation of hydroxyl groups [20]. The main weight loss is observed between 180 and 400°C (denoted as step II), accompanying with a strong exothermic peak, which can be primarily assigned to the decomposition of the surfactant and the combustion of carbon species [21]. Moreover, the weight loss for step I and II of Ce(IV)Cu_x samples are higher than the corresponding step of Ce(III)Cu_x ones with the same copper contents. These results indicate that $(\text{NH}_4)_2\text{Ce}(\text{NO}_3)_6$ precursor can combine with more hydroxyl and CTAB than $\text{Ce}(\text{NO}_3)_3$ precursor during the synthesis process. As a matter of fact, the tetravalent ceric ion formed in the ammonium cerium nitrate aqueous solution




Fig. 2. XRD patterns of CuO–CeO₂ catalysts with different Cu contents: (a) Ce(III)Cux; (b) Ce(IV)Cux.

(Ce(NO₃)₆)^{2−} is actually a stable 12-coordinate icosahedron complex on which each NO₃[−] ligand is bounded to Ce⁴⁺ in its chelate mode by two oxygen atoms [22]. Cerium(IV) is, thus, present in solution as an anionic complex, while cerium(III) is present as a hydrolyzed cation [22]. This difference in charge would benefit the proximity of anionic Ce(NO₃)₆^{2−} and the cationic copper (Cu²⁺) as well as positive electric surfactant (CH₃(CH₂)₁₅N⁺(CH₃)₃) through an electrostatic adsorption mechanism. This hypothesis is also supported by XRF results (Table 1) that the actual Cu contents of Ce(IV)Cux samples are higher than those of Ce(III)Cux ones with the same nominal Cu contents.

3.2. Structure characterization (XRD, LRS, N₂ adsorption–desorption, HRTEM)

3.2.1. XRD and LRS results

Fig. 2 shows XRD patterns of Ce(III)Cux and Ce(IV)Cux samples. As shown in Fig. 2(a), no typical diffraction peaks of crystalline CuO can be observed for Ce(III)Cu5 and Ce(III)Cu15 samples, indicating the existence of well dispersion of CuO species on the surface of CeO₂, or/and the incorporation of copper into the CeO₂ lattice [23]. As Cu content reaches 30 and 50 mol%, the XRD peaks of crystalline CuO ($2\theta = 35.5^\circ, 38.7^\circ$) [JCPDS (Card No. 41-0254)] appear




Fig. 3. Raman spectra of CuO–CeO₂ catalysts with different Cu contents: (a) Ce(III)Cux; (b) Ce(IV)Cux.




Fig. 4. N₂ adsorption-desorption isotherms (a1, a2) and the corresponding distribution curves of pore size (b1, b2) of CuO-CeO₂ catalysts with different Cu contents.

and their intensities increase with the Cu contents. In Fig. 2(b), for Cu content below 50 mol%, only the diffraction peaks of cubic CeO₂ with fluorite structure [JCPDS (Card No. 34-0394)] can be observed for Ce(IV)Cu_x samples. Once Cu content increases to 50 mol%, the characteristic peaks of crystalline CuO appear.

The grain size and lattice parameter of the oxides before and after copper incorporation are summarized in Table 1. The grain size of Ce(III)Cu_x is bigger than that of Ce(IV)Cu_x with the same Cu contents, and all of them gradually decrease with the introduction of more amount of copper species. The lattice parameters of cubic CeO₂ slightly decrease after the addition of copper species,

which provides evidence that some Cu²⁺ ions are incorporated into ceria lattice, probably forming Cu-O-Ce species, by considering that the radius of Cu²⁺ ion (0.073 nm) is smaller than that of Ce³⁺ (0.110 nm) or Ce⁴⁺ ion (0.097 nm) [24]. Additionally, the lattice parameter observed in some case of these samples appears higher than that typically observed for pure CeO₂ (0.5409 nm) [25], which is possibly due to the higher content of Ce³⁺ in these samples than in the reported one, as confirmed by XPS results discussed below.

Raman spectra of the studied samples are shown in Fig. 3. Ce(III)Cu0 and Ce(IV)Cu0 samples exhibit the characteristic F_{2g} vibration band at 460 cm⁻¹ [26]. Once copper species is introduced,




Fig. 5. TEM images of (a1) Ce(III)Cu0, (a2) Ce(III)Cu5, (a3) Ce(III)Cu15, (a4) Ce(III)Cu30, (a5) Ce(III)Cu50, (b1) Ce(IV)Cu0, (b2) Ce(IV)Cu5, (b3) Ce(IV)Cu15, (b4) Ce(IV)Cu30 and (b5) Ce(IV)Cu50.




Fig. 6. HRTEM images of (a1) Ce(III)Cu0, (a2) Ce(III)Cu5, (a3) Ce(III)Cu15, (a4) Ce(III)Cu30, (a5) Ce(III)Cu50, (b1) Ce(IV)Cu0, (b2) Ce(IV)Cu5, (b3) Ce(IV)Cu15, (b4) Ce(IV)Cu30 and (b5) Ce(IV)Cu50.

this peak shifts from 460 to 447 cm^{-1} , and becomes broader and weaker. This shift may imply that changes in the lattice parameter with particle size have occurred, as it was previously reported that a change of the particle size of CeO_2 from 6 nm to 5 μm led to a shift in peak position about 10 cm^{-1} , and was well explained by phonon confinement model [27,28]. Another reason of shifting and broadening may be the presence of oxygen vacancies [29], which is related to structural defects derived from partially incorporation of copper into CeO_2 lattice, in agreement with the decrease of lattice parameter. Additionally, two weak bands at ca. 270 and 590 cm^{-1} are attributed to oxygen vacancies, and the relative concentration of oxygen vacancies can be reflected by the ratio of A_{590}/A_{460} [14] (Table 1). A_{590}/A_{460} values of CuO-CeO_2 are higher than pure CeO_2 ,

indicating the formation of more oxygen vacancies with the incorporation of copper. Moreover, A_{590}/A_{460} values of Ce(III)Cu_x are higher than those of Ce(IV)Cu_x with the same Cu contents, which is related to a higher concentration of oxygen vacancies in Ce(III)Cu_x samples. Thus, as implied from XRD and Raman results, Ce(IV) precursor promotes the formation of CuO-CeO_2 with smaller grain size, while Ce(III) precursor is beneficial to obtain CuO-CeO_2 with higher concentration of oxygen vacancies.

3.2.2. N_2 adsorption–desorption and TEM results

As shown in Fig. 4(a1) and (a2), all samples exhibit IV-type isotherms and H2-type hysteresis loops according to the IUPAC classification, typical for wormhole-like mesostructure and




Fig. 7. XPS of CuO-CeO_2 catalysts: (a), (b) O 1s; (c), (d) Ce 3d; (e), (f) Cu 2p; (g), (h) Cu LMM.




Fig. 7. (continued).

heterogeneously distributed mesoporous radii. The pore size distributions of Ce(III)Cu_x ($x \leq 30$) samples (5–30 nm) are much broader than those of Ce(IV)Cu_x ($x \leq 30$) ones (3–5 nm) (Fig. 4(b1) and (b2)). Ce(III)Cu50 and Ce(IV)Cu50 own pore size distributions in a wide range probably due to the destruct of pore structure by excessive crystalline CuO [23]. Additionally, Ce(IV)Cu_x samples exhibit higher BET surface area and smaller BJH pore size than those of Ce(III)Cu_x with the same Cu contents (Table 1).

Wormhole-like mesostructure, aggregated by metal oxide nanoparticles, can be directly observed for all samples in Fig. 5. The particle sizes of Ce(IV)Cu_x are around 3–4 nm, which are smaller than those of Ce(III)Cu_x (5–10 nm). Structural details of nanoparticles were obtained by HRTEM images in Fig. 6. All samples have the

spacing of lattice fringes around 0.297–0.305 nm, ascribed to the (1 1 1) crystallographic planes of CeO₂. In addition, amorphous surface CuO_x, which are distributed on the top or at the edge of ceria crystal [30], are found for all CuO–CeO₂ samples. Besides, a few CuO crystallites with the lattice spacing 0.225–0.230 nm appear in Ce(III)Cu30, Ce(III)Cu50 and Ce(IV)50 samples. Combined with XRD and Raman results, CuO–CeO₂ catalysts with the lower copper oxide amount (≤ 15 mol% for Ce(III)Cu_x and ≤ 30 mol% for Ce(IV)Cu_x) contain two kinds of copper species: high dispersed CuO species and copper partially incorporation into CeO₂ lattice. Furthermore, catalysts with excess copper oxide (Ce(III)Cu30, Ce(III)Cu50 and Ce(IV)50 samples) include the third copper species: crystalline CuO.




Fig. 8. (a) The relative area of the u_0 (v_0) and u' (v') peaks to the area of Ce 3d region (Ce(III)%) for CuO–CeO₂ catalysts; (b) the ratio of the intensities of Cu 2p satellite peaks to those of the principal peaks (I_{sat}/I_{pp}) for CuO–CeO₂ catalysts; (c) the ratio of Cu⁺ peak area to the total area (Cu(I)%) for CuO–CeO₂ catalysts.

3.3. Chemical states of CuO–CeO₂ catalysts (XPS results)

XPS was performed in order to further illuminate the surface composition and the chemical state of the elements existing in CuO–CeO₂ catalysts. The surface atomic ratios of Cu/Ce (except Ce(III)Cu50 and Ce(IV)Cu50) determined by XPS are much higher than the corresponding Cu/Ce ratios determined by XRF (Table 1), indicating the enrichment of copper species on the catalysts surface. Fig. 7(a and b) shows the O1s spectra of CuO–CeO₂ catalysts. In addition to the presence of the main peak O' at 529.3 eV attributed to the lattice oxygen of the metal oxides, an apparent shoulder peak O'' at 531.2 eV representing the absorbed oxygen in hydroxyl and carbonate groups is observed [29].

The complex spectrum of Ce 3d is decomposed into ten components with the assignment defined in Fig. 7(c and d). The two sets of spin–orbital multiplets, corresponding to the 3d_{3/2} and 3d_{5/2} contributions, are labeled as u and v , respectively. A favorable method, which considers the relative area of the u_0 (v_0) and u' (v') peaks to the area of Ce 3d region, is applied to estimate Ce³⁺ content [31]:

$$\text{Ce(III)\%} = \frac{100 \times [S(u_0) + S(u') + S(v_0) + S(v')]}{\sum [S(u) + S(v)]} \%$$

The Ce(III)% values for the two series of CuO–CeO₂ catalysts are reported in Fig. 8(a). It can be seen that there is a certain amount of Ce³⁺ in all the catalysts, and that the Ce³⁺ quantities of Ce(III)Cux samples are slightly higher than those of Ce(IV)Cux ones with the same Cu contents. Additionally, the values of Ce(III)% for these samples are much higher than that of pure CeO₂ reported in the literature (Ce(III)% = 6.75%) [25], which leads to the higher lattice parameters of these samples than that of the reported one.

Cu 2p spectra in Fig. 7(e and f) shows the principal peaks of Cu 2p_{3/2} at about 934.0 eV and corresponding satellite peaks at 938–948 eV with shake-up structure. Additionally, a weak Cu 2p_{3/2} peak centered at 932.4 eV can be observed. It is reported that the higher Cu 2p_{3/2} binding energy and the shake-up peak are two major XPS characteristics of Cu²⁺ species, while the lower Cu 2p_{3/2} binding energy and absence of the shake-up peak are characteristic of reduced copper species [32]. Thus, most Cu species initially present in these samples are in divalent oxidation state and a

small amount in lower-valence state (Cu⁺ and/or Cu⁰). The reduced degree of Cu species can be investigated by determining the ratio of the intensities of the satellite peaks to those of the principal peaks (I_{sat}/I_{pp} , Fig. 8(b)) [10]. Due to the photoreduction in the spectrometer, it is difficult to estimate the accurate reduction degree of Cu species, but this method gives valuable information for comparison between these two series samples [10]. After the comparison between bulk CuO (I_{sat}/I_{pp} = 0.57) and these samples, a slightly reduced copper phase is observed for all samples. Moreover, the I_{sat}/I_{pp} values for Ce(III)Cux are lower than those for Ce(IV)Cux, indicating the appearance of more amount of low-valence copper species in Ce(III)Cux samples. In order to clarify the low valence state of copper species, Auger Cu LMM lines were also investigated over these mentioned catalysts, as shown in Fig. 7(g and h). It can be observed that the reduced copper species mainly exist as Cu⁺ (916.0 eV) [33]. As shown in Fig. 8(c), the ratios of Cu⁺ area to the total area (denoted as Cu(I)%) for Ce(III)Cux samples are higher than those for Ce(IV)Cux, suggesting more amount of Cu⁺ species in Ce(III)Cux samples.

As discussed above, Ce(III)Cux samples contain more amount of Ce³⁺ (Ce(III)%) and Cu⁺ (I_{sat}/I_{pp} , Cu(I)%) species than Ce(IV)Cux ones, which are expected to exhibit different reduction properties and CO oxidation activities subsequently. Nevertheless, there is more information that can be obtained from Fig. 8(a and b). The ratios of Ce³⁺ and Cu²⁺ in CuO–CeO₂ catalysts decrease simultaneously as more amount of Cu species incorporation. These results provide unambiguous evidence of existence of redox equilibrium of Cu²⁺ + Ce³⁺ \leftrightarrow Cu⁺ + Ce⁴⁺, which has been claimed to be the source of a synergistic interaction between copper and ceria species in the CuO–CeO₂ catalysts. Furthermore, the ratios of Ce³⁺ and Cu²⁺ in Ce(III)Cux samples decline in a higher degree than those in Ce(IV)Cux ones, which indicates that the redox cycles between Cu²⁺/Cu⁺ and Ce³⁺/Ce⁴⁺ are more facile to occur in Ce(III)Cux than in Ce(IV)Cux.

3.4. Reduction properties of CuO–CeO₂ catalysts (H₂-TPR, in situ CO-IR)

3.4.1. H₂-TPR results

Fig. 9(a) displays the H₂-TPR profiles of Ce(III)Cux samples. One peak appears at 491 °C in Ce(III)Cu0, belonging to the reduction of surface oxygen species [15], and the other peak at higher temperature is not included here. Ce(III)Cu5 and Ce(III)Cu15 exhibit two reductive peaks (denoted as α and β), which are attributed to the reduction of well dispersed copper species and copper incorporated into CeO₂ lattice, respectively [23]. For Ce(III)Cu50 sample, a new reduction peak γ at about 185 °C appears, corresponding to the reduction of crystalline CuO [34]. For Ce(III)Cu30 sample, due to the appearance of a small amount of crystalline CuO detected in XRD pattern, the reduction peak at 168 °C is considered partially from the contribution of crystalline CuO. Accordingly, this peak is nominated as peak δ , which is the overlap of peak β and γ . With the increase of Cu content, the temperatures of peak α decrease from 181 °C to 138 °C, and for peak β the temperatures decrease from 217 °C to 164 °C, which is probably due to that the increase of Cu–O–Ce species with Cu contents facilitates the reduction of copper species by considering the reducibility of Cu–O–Ce species [35]. Furthermore, H₂ consumption during the reduction still surpasses the one needed for complete Cu²⁺ to Cu⁰ reduction (Table 1), which is associated with the concurrent reduction of copper species and CeO₂ surface oxygen. As pointed out by Martínez-Arias et al. [10,36], CeO₂ reduction can enhance the reducibility of dispersed CuO. Thus, the reduction of cerium entity promotes the reduction of copper species, leading to lower reduction temperature. Besides, for Ce(III)Cu50 sample the actual amount of H₂ consumption is




Fig. 9. H_2 -TPR profiles of $CuO-CeO_2$ catalysts with different Cu contents: (a) $Ce(III)Cu_x$; (b) $Ce(IV)Cu_x$.

approximate to the theoretical amount of H_2 consumption, which implies that the reduction of CeO_2 surface oxygen species is hindered due to the covering from excess crystalline CuO .

Fig. 9(b) is the H_2 -TPR profiles of $Ce(IV)Cu_x$ samples. For $Ce(IV)Cu_0$ sample, the peak at $487^\circ C$ is ascribed to the reduction of CeO_2 . When Cu contents are 5, 15 and 30 mol%, peak α and β can be observed. For $Ce(IV)Cu_{50}$ sample, the reduction peak δ at $184^\circ C$ appears. Peak α moves from $200^\circ C$ to $155^\circ C$ in $Ce(IV)Cu_5$ and $Ce(IV)Cu_{15}$, and keeps at $155^\circ C$ even when Cu content increases continually. Meanwhile, peak β shifts to lower temperature gradually with the increase of Cu content. The actual amount of H_2 consumption is still larger than the theoretical amount of H_2 consumption for this series of $CuO-CeO_2$ samples (Table 1), which indicates the reduction of surface oxygen species of ceria is involved. Additionally, the actual amount of H_2 consumption gradually approaches the theoretical amount of H_2 consumption with high Cu content ($Ce(IV)Cu_{50}$), indicating the block of the surface oxygen species of ceria from crystalline CuO .

Compared H_2 -TPR results of the two series of crystals, it can be observed that the reduction temperatures of α and β for $Ce(III)Cu_x$ samples are lower than those for $Ce(IV)Cu_x$ samples with the same Cu contents, implying that the reduction occurs more easily in $Ce(III)Cu_x$ than in $Ce(IV)Cu_x$. As demonstrated by XPS results, the redox cycles between Cu^{2+}/Cu^+ and Ce^{3+}/Ce^{4+} are more facile to occur in $Ce(III)Cu_x$ than in $Ce(IV)Cu_x$ due to more Ce^{3+} in $Ce(III)Cu_x$, which subsequently brings a stronger synergistic interaction between copper and ceria species in $Ce(III)Cu_x$. According to reference [37], the strong synergistic effect between copper and cerium can facilitate the reduction of $CuO-CeO_2$. Thus, the reduction behavior of $Ce(III)Cu_x$ is improved greatly.

3.4.2. In situ CO-IR spectra results

To further gain insight into the reduction property of the catalysts, in situ FT-IR of CO adsorption was measured in a TPR procedure, and the results have been shown in Fig. 10. Both of the samples display bands in the highest wavenumber zone corresponding mainly to hydroxyl species. These bands at 3710 , 3636

and 3522 cm^{-1} are attributed to isolated hydroxyls; additionally, associated hydroxyl species giving a broad band extending from ca. 3800 to 2800 cm^{-1} are detected [38,39]. For $Ce(III)Cu_{15}$ sample (Fig. 10(a)), the IR band at 2107 cm^{-1} , attributed to the linear CO adsorbed on Cu^+ sites [40,41], increases gradually from $20^\circ C$, reaches the maximum at $100^\circ C$, and eventually vanishes at $220^\circ C$. The band centered at 2360 cm^{-1} for gaseous CO_2 appears at $80^\circ C$, suggesting the incoming CO reduces the surface species of oxides. The bands at 2940 , 2845 , 1510 and 1386 cm^{-1} are different mode assignments of formate, i.e., ν_{O-C-O} (asym) + δ_{C-H} , ν_{C-H} , ν_{O-C-O} (asym) and ν_{O-C-O} (sym) [38,42]. The bands at 1306 , 1475 , 1359 and 1062 cm^{-1} , ascribed to bi- and mono- or poly dentate carbonate [38], are independent of the temperature. Regarding $Ce(IV)Cu_{15}$ sample (Fig. 10(b)), Cu^+-CO species at 2107 cm^{-1} and gaseous CO_2 at 2360 cm^{-1} appear simultaneously at $20^\circ C$, and the intensity of IR band for Cu^+-CO shows a maximum at $120^\circ C$ and a disappearance at $200^\circ C$. Bi- and mono- or poly dentate carbonates, as well as formate species, are also detected at various temperatures from 20 to $240^\circ C$. While the bands at 1615 , 1391 and 1045 cm^{-1} are considered as hydrogen-carbonate species [38,43], which are not observed in the IR spectra of $Ce(III)Cu_{15}$ sample.

It can be noted that there are some differences between $Ce(III)Cu_{15}$ and $Ce(IV)Cu_{15}$ catalysts in FT-IR spectra: (1) Cu^+ species can be probed by CO at 20 – $220^\circ C$ in $Ce(III)Cu_{15}$, while in $Ce(IV)Cu_{15}$ Cu^+ can be detected at 20 – $200^\circ C$, indicating that Cu^+ in $Ce(III)Cu_{15}$ is more stable than that in $Ce(IV)Cu_{15}$. As discussed above, the $Ce(III)$ -derived sample, containing more Ce^{3+} , has greater ability to assist the redox equilibrium $Cu^{2+} + Ce^{3+} \leftrightarrow Cu^+ + Ce^{4+}$ in shifting to right to form more stable Cu^+ species, which is more difficult for the $Ce(IV)$ -derived sample. (2) Gaseous CO_2 (2360 cm^{-1}) appears at $80^\circ C$ for $Ce(III)Cu_{15}$ sample, while this species appears at $20^\circ C$ for $Ce(IV)Cu_{15}$. CO_2 formation is due to the interaction of CO with Cu^{2+} ions with unsaturated coordination leading to their reduction, while subsequent CO adsorption on reduced centers produces the $Cu^+-carbonyl$ species [9]. The relation between gaseous CO_2 and Cu^+-CO is revealed in Fig. 11. For $Ce(III)Cu_{15}$ sample, CO_2 is not detected from 20 to $60^\circ C$, and the intensities




Fig. 10. In situ FT-IR spectra of CO adsorption on (a) Ce(III)Cu15; (b) Ce(IV)Cu15.

of $\text{Cu}^+ \text{-CO}$ keep the same. As CO_2 appears at 80 °C, the intensity of $\text{Cu}^+ \text{-CO}$ increases. It means that Cu^+ is contained in Ce(III)Cu15 sample, and can exist steadily at 20–60 °C. For Ce(IV)Cu15 sample, the intensities of CO_2 elevate gradually with the increase of temperature from 20 to 60 °C, and the intensities of $\text{Cu}^+ \text{-CO}$ increase simultaneously. These results indicate that Cu^+ is produced by the reduction of Cu^{2+} in Ce(IV)Cu15 sample, accompanying with CO_2 appearance. This difference accounts for that the stability of Cu^+ is enhanced by electron interaction between $\text{Cu}^{2+}/\text{Cu}^+$ and $\text{Ce}^{3+}/\text{Ce}^{4+}$ in Ce(III)-derived sample. (3) Hydrogen-carbonate species can be detected at 1615, 1391 and 1045 cm^{-1} for Ce(IV)Cu15, which are very stable even after heat treatment up to 240 °C. However, these carbonates are not observed for Ce(III)Cu15. Carbonates absorbed on the catalyst surface can prevent the $\text{Ce}^{3+}/\text{Ce}^{4+}$ redox interplay at

the copper–cerium interface and poison active sites of the catalyst [44,45], which is adverse for the activity.

3.5. Catalytic tests of CO oxidation

Fig. 12 shows turnover frequency (TOF, s^{-1}) of CO over per copper atom as a function of temperature for CO oxidation reaction over these CuO-CeO_2 catalysts. For Ce(III)Cu x samples, the TOF follows the order of Ce(III)Cu15 > Ce(III)Cu5 > Ce(III)Cu30 > Ce(III)Cu50. For Ce(IV)Cu x ones, the order is ranked as Ce(IV)Cu15 > Ce(IV)Cu30 > Ce(IV)Cu5 > Ce(IV)Cu50. The catalysts




Fig. 11. Integrated intensity of IR spectra for $\text{Cu}^+ \text{-CO}$ and gaseous CO_2 over CuO-CeO_2 catalysts as a function of temperature.




Fig. 12. Turnover frequency (TOF, s^{-1}) of CO over per copper atom in these CuO-CeO_2 catalysts as a function of temperature.





Fig. 13. In situ FT-IR spectra of CO and O₂ co-adsorption on (a) Ce(III)Cu15; (b) Ce(IV)Cu15.

without detected crystalline CuO display better performance than the catalysts containing crystalline CuO, due to the suppression of surface active species by crystalline CuO [10]. According to the results of structure characterization, Ce(IV)Cu_x catalysts possess higher surface area and smaller pore size than Ce(III)Cu_x ones, but the catalytic activities of Ce(IV)Cu_x are not improved obviously compared with the corresponding Ce(III)Cu_x ones. In this sense, the surface area and the pore size of catalysts are not the main factors influencing the activity. On the other hand, Ce(III)Cu_x catalysts exhibit excellent reduction property with low reduction temperatures in H₂-TPR and highly stable Cu⁺ species in CO-IR, which are originated from the enhanced copper–cerium synergic effect due to high concentration of oxygen vacancies and Ce³⁺. As Martínez-Arias et al. [36,46] reported, the reaction mechanism of CO oxidation for CuO–CeO₂ catalyst is proposed to involve reductive and oxidative single steps, and the rate-determining one is attributed to the oxidative process. However, the oxidative process depends on the starting degree of reduction attained by the system. In other words, if the system attains a relatively high reduction degree, oxidation of the system can occur sequentially with the ceria component becoming oxidized most preferentially [7]. Accordingly, the catalyst which is more easily to be reduced can exhibit higher reaction activity. Besides, Ce(III)Cu_x catalysts own stable Cu⁺ species and high concentration of oxygen vacancies, which provide stable sites for CO chemisorption and active oxygen for reaction to enhance activities. Thus, the different synergistic interactions between Cu and Ce leading to different reduction properties are the main factors influencing the CO oxidation activity. According to our previous study [47], the role of synergistic interaction between copper and ceria in this reaction is tentatively proposed as follows (Scheme 1): considering the reducibility of Cu–O–Ce species [35], the surface Cu–O–Ce species could be reduced by CO in the mixed atmosphere to form surface synergetic oxygen vacancy (SSOV) (i.e. Cu⁺–□–Ce³⁺ species), accompanying by an exposed Cu⁺ and a neighboring Ce³⁺ (Step 1), and the exposed Cu⁺ and oxygen vacancy could provide adsorbed site for CO and activated site for O₂, respectively (Step 2). Then the active oxygen reacts facily with the adsorbed CO species to form CO₂ (Step 3).

3.6. CO and O₂ co-adsorption in situ FT-IR spectra results

In order to further probe the surface species and investigate the influencing factors of the CO oxidation activity, CO and O₂ co-adsorption in situ FT-IR spectra of Ce(III)Cu15 and Ce(IV)Cu15 samples were recorded, as shown in Fig. 13. Hydroxyl species can be detected in the range from ca. 4000 to 2800 cm^{−1} [38]. The two rotational branches, centered at ca. 2360 cm^{−1}, and a band at 2142 cm^{−1} are corresponding to gaseous CO₂ and CO, respectively [48]. The band at 2107 cm^{−1} is observed already upon first contact of the copper-containing samples with CO at room temperature and can be ascribed to Cu⁺–carbonyl species in accordance with previous works [40,49]. Similar IR spectra at 1600–1000 cm^{−1} assignable to bi- and mono- or poly dentate carbonates, as well as formate species [38,42], are clearly seen over Ce(III)Cu15 and Ce(IV)Cu15 samples, which seem to be independent of temperature. New bands at ca. 1615, 1391, 1045 and 1010 cm^{−1} for hydrogen–carbonate and bidentate carbonates can be distinguished in the IR spectra of Ce(IV)Cu15 sample [9,39]. These chemisorbed carbonates are very stable even after heating to 150 °C.

More important information can be obtained from the comparison of the stability of copper carbonyl and adsorbed carbonate species. It is well recognized that the chemisorption of CO on copper species is important for CO oxidation reaction [14]. Although



Scheme 1. The proposed reaction mechanism of CO oxidation over CuO–CeO₂ catalysts.

the band for $\text{Cu}^+ - \text{CO}$ appears at room temperature for both of the samples, this band intensity of Ce(III)Cu15 is much higher than that of Ce(IV)Cu15 . The respective intensities of these $\text{Cu}^+ - \text{carbonyls}$ provide a measure of the potential of each catalyst to become interfacially reduced upon interaction of CO with its copper oxide entities, whereas the level of reduction attained in each case is in turn related to the amount of sites available for oxygen to react and close the catalytic cycle [50]. In this sense, the CO oxidation activity is proportional to the intensity of the $\text{Cu}^+ - \text{carbonyl}$ band. It is consistent with the results of catalytic tests that Ce(III)Cu15 exhibits higher activity than Ce(IV)Cu15 . Moreover, the feature of $\text{Cu}^+ - \text{CO}$ disappears at 110°C for Ce(IV)Cu15 sample, while for Ce(III)Cu15 sample, the desorption process is very slow and this feature still remains at 150°C . The difference in decomposition of $\text{Cu}^+ - \text{CO}$ species between Ce(III)Cu15 and Ce(IV)Cu15 implies that $\text{CuO} - \text{CeO}_2$ catalyst obtained from trivalent ceric precursor has the potential to facilitate the redox equilibrium of $\text{Cu}^{2+} + \text{Ce}^{3+} \leftrightarrow \text{Cu}^+ + \text{Ce}^{4+}$ shifting to right and the stabilization of Cu^+ on the surface, which is beneficial for the enhancement of CO oxidation activity. Apart from the above discussion, it is found that the adsorbed hydrogen-carbonate species over Ce(IV)Cu15 , which are not observed on Ce(III)Cu15 , are difficult to decompose. Active sites can be blocked by carbonate species, resulting in surface poisoning [49]. It is reasonable that the carbonate species would limit reaction rates and lead to low activity of Ce(IV)Cu15 [51].

4. Conclusions

As demonstrated by comprehensive characterizations, the use of different cerium precursors leads to different properties of mesoporous $\text{CuO} - \text{CeO}_2$ catalysts: (1) the use of $(\text{NH}_4)_2\text{Ce}(\text{NO}_3)_6$ precursor can improve the surface area, pore size and particle size, instead of elevating the reduction property and catalytic activity. (2) The use of $\text{Ce}(\text{NO}_3)_3$ precursor can promote the redox equilibrium of $\text{Cu}^{2+} + \text{Ce}^{3+} \leftrightarrow \text{Cu}^+ + \text{Ce}^{4+}$ shifting to right and the formation of high amount of Cu^+ species and oxygen vacancies, which are responsible for the improved reducibility and activity for CO oxidation. (3) Ce(III) -derived samples contain high amount of Ce^{3+} , leading to facile $\text{Ce}^{4+}/\text{Ce}^{3+}$ redox cycle and meanwhile promotion of copper species reduction. (4) Ce(IV) -derived samples are prone to adsorb carbonate species, which is adverse for the activity of CO oxidation.

Acknowledgments

The financial supports of the National Nature Science Foundation of China (Nos. 20873060, 20973091) and the National 973 Program of China (No. 2010CB732300) are gratefully acknowledged.

References

- [1] M. Kang, M.W. Song, C.H. Lee, Appl. Catal. A: Gen. 251 (2003) 143–156.
- [2] X.L. Tang, B.C. Zhang, Y. Li, Y.D. Xu, Q. Xin, W.J. Shen, Appl. Catal. A: Gen. 288 (2005) 116–125.
- [3] H.Q. Zhu, Z.F. Qin, W.J. Shan, W.J. Shen, J.G. Wang, J. Catal. 225 (2004) 267–277.
- [4] F. Mariño, C. Descorme, D. Duprez, Appl. Catal. B: Environ. 54 (2004) 59–66.
- [5] Q. Fu, A. Weber, M. Flytzani-Stephanopoulos, Catal. Lett. 77 (2001) 87–95.
- [6] J.B. Wang, D.H. Tsai, T.J. Huang, J. Catal. 208 (2002) 370–380.
- [7] A. Martínez-Arias, A.B. Hungria, M. Fernández-García, J.C. Conesa, G. Munuera, J. Phys. Chem. B 108 (2004) 17983–17991.
- [8] W. Liu, M. Flytzani-Stephanopoulos, J. Catal. 153 (1995) 304–316.
- [9] A. Martínez-Arias, M. Fernández-García, J. Soria, J.C. Conesa, J. Catal. 182 (1999) 367–377.
- [10] B. Skärman, D. Grandjean, R.E. Benfield, A. Hinz, A. Andersson, L.R. Wallenberg, J. Catal. 211 (2002) 119–133.
- [11] A. Martínez-Arias, A.B. Hungria, G. Munuera, D. Gamarra, Appl. Catal. B: Environ. 65 (2006) 207–216.
- [12] G. Sedmak, S. Hocevar, J. Levec, J. Catal. 222 (2004) 87–99.
- [13] H.B. Zou, X.F. Dong, W.M. Lin, Appl. Surf. Sci. 253 (2006) 2893–2898.
- [14] Z.Y. Pu, X.S. Liu, A.P. Jia, Y.L. Xie, J.Q. Lu, M.F. Luo, J. Phys. Chem. C 112 (2008) 15045–15051.
- [15] M.F. Luo, Y.P. Song, J.Q. Lu, X.Y. Wang, Z.Y. Pu, J. Phys. Chem. C 111 (2007) 12686–12692.
- [16] W.H. Shen, X.P. Dong, Y.F. Zhu, H.R. Chen, J.L. Shi, Microporous Mesoporous Mater. 85 (2005) 157–162.
- [17] S. Bernal, J.J. Calvino, M.A. Cauqui, J.M. Gatica, C. Larese, J.A. Pérez Omil, J.M. Pintado, Catal. Today 50 (1999) 175–206.
- [18] X.L. Tang, B.C. Zhang, Y. Li, Y.D. Xu, Q. Xin, W.J. Shen, Catal. Today 93–95 (2004) 191–198.
- [19] Y. Li, Q. Fu, M. Flytzani-Stephanopoulos, Appl. Catal. B: Environ. 27 (2000) 179–191.
- [20] B.M. Reddy, P. Lakshmanan, A. Khan, S. Lorient, C. López-Cartes, T.C. Rojas, A. Fernández, J. Phys. Chem. B 109 (2005) 13545–13552.
- [21] J.L. Cao, Y. Wang, X.L. Yu, S.R. Wang, S.H. Wu, Z.Y. Yuan, Appl. Catal. B: Environ. 79 (2008) 26–34.
- [22] S. Letichevsky, C.A. Tellez, R.R. de Avillez, M.I.P. da Silva, M.A. Fraga, L.G. Appel, Appl. Catal. B: Environ. 58 (2005) 203–210.
- [23] M.F. Luo, J.M. Ma, J.Q. Lu, Y.P. Song, Y.J. Wang, J. Catal. 246 (2007) 52–59.
- [24] W.J. Shan, W.J. Shen, C. Li, Chem. Mater. 15 (2003) 4761–4767.
- [25] W.D. Cai, F. Chen, X.X. Shen, L.J. Chen, J.L. Zhang, Appl. Catal. B: Environ. 101 (2010) 160–168.
- [26] R. Kydd, W.Y. Teoh, K. Wong, Y. Wang, J. Scott, Q.H. Zeng, A.B. Yu, J. Zou, R. Amal, Adv. Funct. Mater. 19 (2009) 369–377.
- [27] J.E. Spanier, R.D. Robinson, F. Zhang, S.W. Chan, I.P. Herman, Phys. Rev. B 64 (2001) 245407–245411.
- [28] M. Fernández-García, A. Martínez-Arias, J.C. Hanson, J.A. Rodriguez, Chem. Rev. 104 (2004) 4063–4104.
- [29] L. Ilieva, G. Pantaleo, I. Ivanov, A.M. Venezia, D. Andreeva, Appl. Catal. B: Environ. 65 (2006) 101–109.
- [30] L.J. Liu, Z.J. Yao, B. Liu, L. Dong, J. Catal. 275 (2010) 45–60.
- [31] J. Silvestre-Albero, F. Rodríguez-Reinos, A. Sepúlveda-Escribano, J. Catal. 210 (2002) 127–136.
- [32] B. Wen, M.Y. He, Appl. Catal. B: Environ. 37 (2002) 75–82.
- [33] I. Platzman, R. Brenner, H. Haick, R. Tannenbaum, J. Phys. Chem. C 112 (2008) 1101–1108.
- [34] D. Delimaris, T. Ioannides, Appl. Catal. B: Environ. 89 (2009) 295–302.
- [35] P. Bera, K.R. Priolkar, P.R. Sarode, M.S. Hegde, S. Emura, R. Kumashiro, N.P. Lalla, Chem. Mater. 14 (2002) 3591–3601.
- [36] A. Martínez-Arias, M. Fernández-García, O. Gálvez, J.M. Coronado, J.A. Anderson, J.C. Conesa, J. Soria, G. Munuera, J. Catal. 195 (2000) 207–216.
- [37] W.J. Shan, Z.C. Feng, Z.L. Li, J. Zhang, W.J. Shen, C. Li, J. Catal. 228 (2004) 206–217.
- [38] P. Bera, A.L. Cámara, A. Hornés, A. Martínez-Arias, J. Phys. Chem. C 113 (2009) 10689–10695.
- [39] A. Holmgren, B. Andersson, D. Duprez, Appl. Catal. B: Environ. 22 (1999) 215–230.
- [40] H.Q. Wan, Z. Wang, J. Zhu, X.W. Li, B. Liu, F. Gao, L. Dong, Y. Chen, Appl. Catal. B: Environ. 79 (2008) 254–261.
- [41] O. Dulacourt, X. Courtois, V. Perrichon, D. Bianchi, J. Phys. Chem. B 104 (2000) 6001–6011.
- [42] A. Hornés, P. Bera, A.L. Cámara, D. Gamarra, G. Munuera, A. Martínez-Arias, J. Catal. 268 (2009) 367–375.
- [43] C. Binet, M. Daturi, J.C. Lavalley, Catal. Today 50 (1999) 207–225.
- [44] C.V. Ovensen, P. Stolze, J.K. Nørskov, C.T. Campbell, J. Catal. 134 (1992) 445–468.
- [45] C.H. Kim, L.T. Thompson, J. Catal. 230 (2005) 66–74.
- [46] K. Qian, W.X. Huang, J. Fang, S.S. Lv, B. He, Z.Q. Jiang, S.Q. Wei, J. Catal. 255 (2008) 269–278.
- [47] D. Li, Q. Yu, S.S. Li, H.Q. Wan, L.J. Liu, L. Qi, B. Liu, F. Gao, L. Dong, Y. Chen, Chem. Eur. J. 17 (2011) 5668–5679.
- [48] F. Morales, E. de Smit, F.M.F. de Groot, T. Visser, B.M. Weckhuysen, J. Catal. 246 (2007) 91–99.
- [49] F.C. Meunier, D. Tibiletti, A. Goguet, D. Reid, R. Burch, Appl. Catal. A: Gen. 289 (2005) 104–112.
- [50] D. Gamarra, C. Belver, M. Fernández-García, A. Martínez-Arias, J. Am. Chem. Soc. 129 (2007) 12064–12065.
- [51] S. Hilaire, X. Wang, T. Luo, R.J. Gorte, J. Wagner, Appl. Catal. A: Gen. 215 (2001) 271–278.